

## Immunology Immunoglobulin Referral Form

Phone: **1-877-342-9352**Fax: 1-888-594-4844

>< Please detach before submitting to a pharmacy – tear here.

PATIENT INF	ORMATION												
IG specialist	Name:	Prima	Primary diagnosis:										
	Phone:			☐ Combined immunodeficiency									
Patient	☐ see attached	nale 🔲 Cor	☐ Common hypogammaglobulinemia										
Patient name:_		🗖 Cor	☐ Common variable immunodeficiency										
DOB:			🗖 lmm	_ ☐ Immune thrombocyopenic purpura									
Address:			🗖 lmm	_ ☐ Immunodeficiency with increased IgM									
City:	City:          State:          Zip:			☐ Wiskott-Aldrich syndrome									
Phone: Cell:				_ Other:									
Emergency contact:				_ Medical assessment									
Phone: Relationship:			Height	Height: Weight: O lbs O kg									
Insurance	☐ Front and bac	v Curren	Current medications? O Yes O No										
	Primary	Secondary	y If yes,	If yes, list or attach:									
Insurance:													
Phone:				Allergies:									
Policy #:													
Group:													
	ON ORDERS												
Immune Globu			Initial:		gm/kg	divided over		days					
□ No preference				ng:		divided over							
□ Preferred product:						or							
Directions:				every weeks for cycles  Quantity/Refills:									
						nths unless other	wise no	oted					
<ul><li>□ Infuse IV □ Infuse SC</li><li>☑ Per manufacturer guidelines or as written below:</li></ul>				☐ 1-month supply; refill x 12 months unless otherwise noted ☐ Other:									
				☑ Infusion reaction management and kit order									
☑ May round to the nearest 5gm vial size				Mild Moderate	Slow infusion rate by 50% until symptoms resolve Diphenhydramine PO O 25 mg O 50 mg O other:  Stop infusion, resume at 50% when symptoms resolve Diphenhydramine IV O 25 mg O 50 mg O other:								
Pre-medications 30 minutes before start of IG:													
☐ Acetaminophen PO													
									□ Other:				
									Nursing and other orders:				
☑ Administer IVIG or teach SCIG self-administration, via pump									Administer epinephrine 1 mg/mL by weight (Wt):				
<ul> <li>☑ Ambulatory pump if required for infusion</li> <li>☑ Initiate access device (insert peripheral IV, SC needles, access implanted port, or use existing PICC)</li> </ul>			ss Se	Severe	Wt > 66 lbs			Vt < 33 lbs					
			(Anap	hylaxis)	(30 kg) =	(15 - 30 kg) =	(	<15 kg) =					
☑ Flush PIV with 5mL NS (for other orders, contact physician)			l	II 911*	0.3 mg/.3mL	0.15 mg/0.15m	IL U	.01 mg/kg					
□ Obtain labs (list):					Repeat epinephrine in 5-15 min if symptoms continue.								
Lab frequency: ☐ Once ☐ Monthly ☐ Other:					Administer CPR if needed until EMS arrives.								
PHYSICIAN I	NFORMATION												
Name:		Addr	ress:										
						Zip:							
		NPI:		orm: !!		a ralacca the above referen							
		lucts and services are medically necessary a has my permission to contact the insurance				release line above referend	æu intormati	ion and medical and					
Substitution permissible Signature:Date:				Dispense as written Signature:									